

The quark-antiquark potential at finite temperature and the dimension two gluon condensate

E. Megías,^{*} E. Ruiz Arriola,[†] and L.L. Salcedo
*Departamento de Física Atómica, Molecular y Nuclear,
 Universidad de Granada, E-18071 Granada, Spain*

(Dated: February 2, 2008)

A recently proposed phenomenological model, which includes non perturbative effects from dimension two gluon condensates, is applied to analyze the available lattice data for the heavy quark free energy in the deconfined phase of quenched QCD. For large $q\bar{q}$ separations we recover previous results for the Polyakov loop, exhibiting unequivocal condensate contributions. For the $q\bar{q}$ potential at finite temperature and finite separation we find that a good overall description of the lattice data can be achieved once the condensate is properly accounted for. In addition, the model predicts a duality between the zero temperature potential as a function of the $q\bar{q}$ separation, on the one hand, and the quark selfenergy as a function of the temperature, on the other, which turns out to be satisfied to a high degree by the lattice data.

PACS numbers: 11.10.Wx, 11.10.Kk, 11.15.-q, 11.10.Jj

Keywords: Polyakov Loop; Nonperturbative Effects; Dimension two Condensate; QCD; Lattice QCD; Deconfinement Phase; Thermal Field Theory

I. INTRODUCTION

The understanding of the physics of the quark gluon plasma [1] has triggered a lot of activity in the past years. (See [2, 3] for accounts of the experimental situation.) As a part of this effort, there have been some recent advances in the study of the inter-quark forces at finite temperature [4, 5]. In these analyses the change in free energy due to the presence of a static quark-antiquark pair separated by a distance r in a thermal bath has often been used. The color screening at high temperature will produce a dissolution of the quarkonium state, and this could be a signal of quark gluon plasma formation [6].

The vacuum expectation value of the Polyakov loop is related to the propagator of a static quark and constitutes a natural order parameter for the deconfinement phase transition of quenched QCD. Due to ultraviolet divergences in the heavy quark selfenergy, the Polyakov loop is subject to multiplicative renormalization (see e.g. [7, 8]). The perturbative computation of the renormalized Polyakov loop has been carried out to next to leading order [9] and only recently a practical systematic procedure has been devised to obtain it non perturbatively in lattice QCD, based on the computation of singlet and octet correlation functions (Wilson lines) [10]. These correlation functions are related to the free energy of a heavy quark-antiquark pair. In the limit of large separation the two Polyakov loops decouple, and the result is essentially the square of one single Polyakov loop. In the opposite limit of separations much smaller than the Debye length the result is temperature independent, so that the $q\bar{q}$ free energy reproduces the known QCD static $q\bar{q}$ potential at

zero temperature, $V_{q\bar{q}}(r)$, [11]. There is an additive ambiguity in the free energy that is fixed at short distances by comparison with this zero temperature quantity. This strongly suggests that there exists a deep connection between the zero temperature $q\bar{q}$ potential and the vacuum expectation value of the Polyakov loop.

The perturbative computation of the $q\bar{q}$ potential has been studied for a long time. At tree-level, $V_{q\bar{q}}(r)$ is a Coulomb potential arising from one-gluon exchange. Up to now there are estimates of the three loop perturbative contribution using several methods [12, 13, 14]. The perturbative series turned out to be very poorly convergent at $r \geq 0.1$ fm and resummation prescriptions lead to large uncertainty due to scheme dependence. Several techniques have been proposed to improve the perturbative series: extraction of the $\mathcal{O}(\Lambda_{\text{QCD}})$ infrared renormalons [15, 16], Operator Product Expansion (OPE) for $r \ll \Lambda_{\text{QCD}}^{-1}$ [17], etc. Non-perturbative methods, such as lattice simulations, and phenomenological potential model analyses of experimental data for the heavy quarkonium spectra, indicate that the $q\bar{q}$ potential can be well approximated by the sum of a Coulomb term plus a linear term that becomes dominant at separations of the order $\sim 1/\Lambda_{\text{QCD}}$ [18, 19]. This non-perturbative linear behavior at large distances is consistent with the quark confinement picture based on the QCD string. For a recent attempt within the holographic QCD approach based on the AdS/CFT correspondence see e.g. [20] and references therein.

On the other hand, at temperatures of the order of the deconfinement phase transition the characteristic Euclidean scale corresponds to $\mu \sim 4\pi T_c \sim 3$ GeV, and one expects the OPE formalism to apply [21]. This approach consists of expanding the Green's functions in inverse power series of p^2 , each term corresponding to a local operator, implying that condensates and power corrections could play a role. This has been verified

^{*}Electronic address: emegias@ugr.es

[†]Electronic address: earriola@ugr.es

in a recent analysis of the renormalized Polyakov lattice data [10], where the existence of non perturbative contributions driven by gluon condensates has been exposed [22]. The first BRST invariant condensate is of dimension two $\langle A_{\mu,a}^2 \rangle$, with $A_{\mu,a}$ the gluon field [23]. It is tempting to develop a model based on the computation of correlations of Polyakov loops to connect these two regimes (zero temperature and large distances), so that the non-perturbative contributions in the Polyakov loop have the same origin as those in the zero temperature $q\bar{q}$ potential, namely, the gluon condensates. This will be one of the issues of the manuscript. More generally, we extend the one-gluon exchange model introduced in [22] for the Polyakov loop, to the description of the singlet heavy quark-antiquark free energy and compare with available lattice data. A preliminary study in this direction was presented in [24].

The paper is organized as follows. In section II we review the observed non-perturbative behavior of the renormalized Polyakov loop just above the deconfinement phase transition, and the importance of thermal power corrections to describe it, thereby justifying the introduction of dimension two condensates in the theory. In section III we study the influence of these condensates on the free energy of the static $q\bar{q}$ pair. In section IV we compare our model with available lattice data and establish a remarkable duality between the zero temperature $q\bar{q}$ potential and the Polyakov loop. Finally, in section V we summarize our points and draw our main conclusions.

II. THE POLYAKOV LOOP

In this section we summarize results derived for the Polyakov loop and the role played by the dimension two condensate. Fuller details are given in [22]. The Polyakov loop or thermal Wilson line is the propagator of a static color source. Physically, the expectation value of the n -point Polyakov loop correlation functions are related to the change in free energy arising from the presence of static quark and antiquark sources in the heat bath. The Polyakov loop is defined by

$$\Omega(\mathbf{x}) = \mathcal{T} \exp \left(ig \int_0^{1/T} dx_0 A_0(\mathbf{x}, x_0) \right), \quad (1)$$

where $\mathcal{T}$ denotes time ordering, and x_0 is the Euclidean time coordinate. $A_0 = \sum_a A_{0,a} T_a$ and T_a , $a = 1, \dots, N_c^2 - 1$, are the generators of $SU(N_c)$ in the fundamental representation, with normalization $\text{tr}(T_a T_b) = \delta_{ab}/2$, and N_c the number of colors. Let

$$L(T) = \left\langle \frac{1}{N_c} \text{tr} \Omega \right\rangle, \quad (2)$$

be the normalized expectation value of the Polyakov loop (tr is the trace in the fundamental representation of the color group). This quantity was computed perturbatively

to next-to-leading order (NLO) by Gava and Jengo [9]. To leading order (LO),

$$L(T) = 1 + \frac{1}{16\pi} \frac{N_c^2 - 1}{N_c} g^2 \hat{m}_D + \mathcal{O}(g^4), \quad (3)$$

where we have defined the dimensionless quantity $\hat{m}_D = m_D/T$, and m_D is the Debye mass. To one loop [25]

$$\hat{m}_D = \frac{m_D}{T} = g(N_c/3 + N_f/6)^{1/2} \quad (4)$$

N_f being the number of flavors. The temperature dependence in the perturbative L comes from the running of the coupling constant.

In a static gauge (i.e., one where A_0 is time independent), the Polyakov loop takes the simpler form

$$\Omega(\mathbf{x}) = e^{igA_0(\mathbf{x})/T}, \quad (5)$$

and the computation of $L(T)$ can be undertaken using dimensional reduction ideas [25, 26, 27, 28, 29]. Namely, after the non stationary Matsubara modes are integrated out, the static modes $A_0(\mathbf{x})$ and $\mathbf{A}(\mathbf{x})$ are described by an effective three dimensional Lagrangian with computable parameters.¹ Expanding the exponential one finds²

$$L(T) = 1 - \frac{g^2}{2T^2} \frac{1}{N_c} \langle \text{tr}(A_0^2) \rangle + \frac{g^4}{24T^4} \frac{1}{N_c} \langle \text{tr}(A_0^4) \rangle + \dots \quad (6)$$

The quartic term is $\mathcal{O}(g^6)$ and hence, to order $\mathcal{O}(g^5)$ one can use the Gaussian-like approximation

$$L = \exp \left[-\frac{g^2 \langle A_{0,a}^2 \rangle}{4N_c T^2} \right] + \mathcal{O}(g^6). \quad (7)$$

This formula becomes exact in the large N_c limit [30]. From here it is immediate to relate the Polyakov loop to the gluon propagator of the dimensionally reduced theory

$$\langle A_{0,a}(\mathbf{x}) A_{0,b}(\mathbf{y}) \rangle = \delta_{ab} T \int \frac{d^3 k}{(2\pi)^3} e^{i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})} D_{00}(\mathbf{k}). \quad (8)$$

The lowest order perturbative term,

$$D_{00}^P(\mathbf{k}) = \frac{1}{\mathbf{k}^2 + m_D^2} + \mathcal{O}(g^2), \quad (9)$$

¹ A subtle point is that to make contact with the Polyakov loop, the integration of non static Matsubara modes should be done in a static gauge, whereas most calculations of the dimensionally reduced Lagrangian are carried out in a covariant gauge. Fortunately the spurious gauge dependence so generated only affects higher perturbative orders [22].

² Odd orders vanish assuming that the conjugation symmetry $A_\mu(x) \rightarrow -A_\mu^T(x)$ is not spontaneously broken. By the same token $\langle \Omega \rangle$ and $\langle \Omega^\dagger \rangle$ are real (and equal in the absence of baryonic chemical potential, which we assume throughout).




FIG. 1: The logarithm of the renormalized Polyakov loop in gluodynamics. Lattice data from [10] for $N_\tau = 4$ (white squares) and $N_\tau = 8$ (black squares). The straight line uses Eq. (11) with the continuum extrapolated values of a and b in (12) (rather than the best fits to the $N_\tau = 4$ or $N_\tau = 8$ data). LO and NLO perturbative calculations [9] are shown for comparison.

(the upperscript P refers to *perturbative* contribution) yields

$$\langle A_{0,a}^2 \rangle^P = -(N_c^2 - 1) \frac{\hat{m}_D}{4\pi} T^2 + \mathcal{O}(g^2), \quad (10)$$

where dimensional regularization rules have been applied to deal with the UV divergence at coincident points. When combined with the Gaussian-like formula (7) this term reproduces to LO the perturbative result in (3). The NLO result is also reproduced in this way [22].

Because in perturbation theory the dimensionless ratio T/Λ_{QCD} appears only through logarithms from radiative corrections, the finite temperature perturbative condensate $\langle A_{0,a}^2 \rangle^P$ is roughly proportional to T^2 and the perturbative $L(T)$ is rather flat as a function of T . This is definitively not what is seen in the lattice data, instead in the unconfined phase one finds the pattern

$$-2 \log L = a + b \left(\frac{T_c}{T} \right)^2 \quad (11)$$

with a and b weakly dependent on T . This is shown in Fig. 1 for gluodynamics data [10]³. The same pattern applies for the unquenched data of [32]. For later reference we quote the estimated continuum limit of the fitted values of a and b for gluodynamics [22]

$$a = -0.19(5), \quad b = 1.63(16). \quad (12)$$

A roughly constant a can be accommodated with perturbation theory, and this is so quantitatively in the high temperature region. The second piece involves a power-like (as opposed to logarithmic) temperature dependence of non perturbative origin, since T_c is proportional to Λ_{QCD} . On purely dimensional grounds it suggests a temperature independent non perturbative dimension two condensate to be added in $\langle A_{0,a}^2 \rangle$. At zero temperature non perturbative condensates appear naturally within the operator product expansion approach [21, 33] through modifications in the propagators of particles with respect to their perturbative counterparts. E.g. the standard dimension four gluon condensate [34, 35] implies a term of the type $\langle G_{\mu\nu}^2 \rangle / \mathbf{k}^6$ to be added to the perturbative gluon propagator $1/\mathbf{k}^2$. Such non perturbative addition is irrelevant in the very high momentum regime but introduces sizeable power-like corrections as one approaches the strong interacting and non perturbative low momentum regime. A dimension two condensate has been proposed in the same zero temperature context in [36, 37]. This implies a gluon propagator of the type $1/\mathbf{k}^2 + m_G^2/\mathbf{k}^4$ which, on dimensional grounds, yields a gluon exchange potential of the form Coulomb plus string tension, $V \sim c/r + \sigma r$ (with σ positive provided m_G^2 is also positive, corresponding to a tachyonic gluon squared mass $-m_G^2$). As we will see immediately, such m_G^2 gives rise to a dimension two condensate in the gluon field.

Guided by this insight the same approach is taken in [22] to describe Polyakov loop data at not so high temperatures above the deconfining transition. That is, a non perturbative piece driven by a dimension two operator is added in the finite temperature propagator:

$$D_{00}(\mathbf{k}) = D_{00}^P(\mathbf{k}) + D_{00}^{\text{NP}}(\mathbf{k}) \quad (13)$$

where

$$D_{00}^{\text{NP}}(\mathbf{k}) = \frac{m_G^2}{(\mathbf{k}^2 + m_D^2)^2}. \quad (14)$$

This expression nicely combines the effects coming from high frequency modes (the Debye mass from integration of the heavy thermal modes) and from the low frequency modes incorporated in the condensates. Thus we expect it to work in a window range above the phase transition.

Using (8) at $\mathbf{x} = \mathbf{y}$, the new term produces

$$\langle A_{0,a}^2 \rangle = \langle A_{0,a}^2 \rangle^P + \langle A_{0,a}^2 \rangle^{\text{NP}} \quad (15)$$

where the non perturbative condensate is related to m_G^2 through

$$\langle A_{0,a}^2 \rangle^{\text{NP}} = \frac{(N_c^2 - 1)m_G^2}{8\pi\hat{m}_D}. \quad (16)$$

This contribution is UV finite. When inserted in (7) it produces the term $b(T_c/T)^2$ in the pattern (11), assuming that, m_G^2 or $\langle A_{0,a}^2 \rangle^{\text{NP}}$ are temperature independent modulo radiative corrections. For future reference we introduce the notation

$$\mathcal{C}_2 := g^2 \langle A_{0,a}^2 \rangle^{\text{NP}} \quad (17)$$

³ An alternative approach is taken in [31] to measure the renormalized Polyakov loop in lattice QCD.

to denote the non perturbative dimension two condensate. Remarkably, the numerical value of the non perturbative finite temperature condensate $\mathcal{C}_2$ turns out to be consistent [22] with independent determinations of the zero temperature condensate [38, 39, 40]. This is in favor of the consistency of the present approach.

III. THE SINGLET $Q\bar{Q}$ FREE ENERGY

In this section we apply the previous model to the study of the free energy of a heavy quark-antiquark pair. The latter is obtained through the correlation of two Polyakov loops,

$$e^{-F_{q\bar{q}}(r,T)/T} = \left\langle \frac{1}{N_c} \text{tr} \Omega(\mathbf{x}) \frac{1}{N_c} \text{tr} \Omega^\dagger(\mathbf{y}) \right\rangle, \quad (18)$$

where $r = |\mathbf{x} - \mathbf{y}|$ is the separation between quark and antiquark. The free energy contains the finite temperature $q\bar{q}$ potential as well as the quarks selfenergies:

$$F_{q\bar{q}}(r, T) = V_{q\bar{q}}(r, T) + 2\Sigma_q(T), \quad (19)$$

where (twice) the selfenergy

$$2\Sigma_q(T) = \lim_{r \rightarrow \infty} F_{q\bar{q}}(r, T) \quad (20)$$

is finite in the deconfined phase and in this case the potential vanishes at large separations. In the confining phase the selfenergy would be infinite and $F_{q\bar{q}}(r, T)$ is directly identified with the $q\bar{q}$ potential $V_{q\bar{q}}(r, T)$. Of course, at large separations the two Polyakov loops in (18) become uncorrelated implying that the quark (or antiquark) selfenergy is related to the expectation value of the Polyakov loop through

$$L(T) = e^{-\Sigma_q(T)/T}. \quad (21)$$

Because the interaction between the color singlets $\text{tr} \Omega$ and $\text{tr} \Omega^\dagger$ requires the exchange of at least two gluons (each carrying a Debye screening factor $e^{-m_D r}$ at large separations), this so called *color averaged* free energy is less suited to lattice computation than the *singlet* free energy

$$e^{-F_1(r,T)/T} = \left\langle \frac{1}{N_c} \text{tr} (\Omega(\mathbf{x}) \Omega^\dagger(\mathbf{y})) \right\rangle. \quad (22)$$

This quantity defined in the Coulomb gauge acquires a gauge invariant meaning [41] and lattice calculations for it are available in the literature [42]. The corresponding $q\bar{q}$ potential is dominated by a single gluon exchange falloff at large separations.

Following precisely the same approach described in the previous section for the Polyakov loop, we obtain the relation analogous to (7)

$$\left\langle \frac{1}{N_c} \text{tr} (\Omega(\mathbf{x}) \Omega^\dagger(\mathbf{y})) \right\rangle = e^{\frac{g^2}{2N_c T^2} (\langle A_{0,a}(\mathbf{x}) A_{0,a}(\mathbf{y}) \rangle - \langle A_{0,a}^2 \rangle)} \quad (23)$$

modulo $\mathcal{O}(g^6)$ corrections. Once again, based on the cumulant expansion [30] these corrections vanish in the large N_c limit. From the relation $F_1 = V_1 + 2\Sigma_q$ similar to (19)⁴ we identify the two contributions

$$\begin{aligned} V_1(r, T) &= -\frac{g^2}{2N_c T} \langle A_{0,a}(\mathbf{x}) A_{0,a}(\mathbf{y}) \rangle, \\ \Sigma_q(T) &= \frac{g^2}{4N_c T} \langle A_{0,a}^2 \rangle, \end{aligned} \quad (24)$$

and in particular the results of the previous section for the Polyakov loop expectation value are recovered with (at LO)

$$\Sigma_q(T) = \frac{g^2(N_c^2 - 1)}{16\pi N_c} \left(-\hat{m}_D T + \frac{m_G^2}{2\hat{m}_D} \frac{1}{T} \right). \quad (25)$$

For the color singlet potential we derive the expression (at LO and neglecting any scale dependence in the coupling constant)

$$V_1(r, T) = -\frac{g^2(N_c^2 - 1)}{8\pi N_c} \left(\frac{1}{r} + \frac{m_G^2}{2\hat{m}_D} \frac{1}{T} \right) e^{-\hat{m}_D T r}. \quad (26)$$

In these results the Polyakov loop is already renormalized by dimensional regularization. In the corresponding lattice computation the bare Polyakov loop has to be renormalized using a suitable prescription. Specifically, in [10, 42] the following criterion is used: the multiplicative renormalization of the Polyakov loop translates into an additive renormalization in the $q\bar{q}$ free energy,

$$F_1^{\text{renorm}}(r, T) = F_1^{\text{bare}}(r, T) + c(T), \quad (27)$$

therefore, because at short distances the renormalized free energy ought to be temperature independent, a suitable r -independent (but T -dependent) constant $c(T)$ is added in such a way that at short distances the (renormalized) finite temperature free energy $F_1(r, T)$ has a maximum overlap with the zero temperature one $F_1(r, 0)$. The contact between the universal zero temperature curve and the temperature dependent one occurs for all separations below a distance of the order of the Debye length [10]. This device allows a quite accurate lattice determination of the selfenergy, and hence of the renormalized Polyakov loop. In practical terms the above criterion means that, if the free energy is expanded in powers of the separation,

$$F_1(r, T) = v_0/r + v_1 + v_2 r + \mathcal{O}(r^2) \quad (28)$$

(neglecting the logarithmic dependence from the running of the coupling constant), the constant term v_1 should be absent (i.e. the Cornell prescription). This is because the

⁴ Due to $\text{SU}(N_c)$ invariance, $\langle \Omega \rangle$ is a multiple of the identity matrix and so Σ_q is the same quantity in $F_{q\bar{q}}$ and F_1 .

dominant Coulombian term v_0/r is temperature independent, and v_1 vanishes in $F_1(r, 0)$ which is of the Coulomb plus string tension form. (A non-vanishing v_1 would require an inexistent “dimension one” condensate.)

As it turns out, our formulas (25) and (26) automatically satisfy this Cornell-Bielefeld prescription for the free energy. I.e., in $F_1 = V_1 + 2\Sigma_q$ the term v_1 is already zero without any further renormalization. This is most remarkable since the selfenergy used a dimensionally regularized definition of the UV divergent integral $\int d^3k D_{00}^P(\mathbf{k})$ leading to (10), whereas the potential is UV finite for any non vanishing value of r . Actually, this was to be expected because, as is well-known, dimensional regularization does not introduce any power-like dependence in addition to those already existing in the theory, so a dimensionful v_1 could not be introduced in the calculation. By the same token we expect that the Cornell-Bielefeld prescription will be automatically satisfied also at higher orders, i.e., beyond the Gaussian-like approximation, provided dimensional regularization is used. Note also that the fact that the Cornell-Bielefeld prescription used in lattice is satisfied by our approach justifies the comparison of our calculation of the renormalized Polyakov loop, based on single loops, with that in lattice, based on the correlation of pairs of loops.

IV. COMPARISON WITH LATTICE DATA

A. Duality between the zero temperature $Q\bar{Q}$ -potential and the Polyakov loop

In this subsection we will analyze within our model the $q\bar{q}$ singlet free energy with respect to the two asymptotic behaviors of zero temperature (at finite separation) and large separation (at finite temperature). Neglecting momentarily any scale dependence in both the coupling constant and the non perturbative condensate, we obtain (at LO)

$$F_1(\infty, T) = -\frac{(N_c^2 - 1)}{8\pi N_c} g^2 \hat{m}_D T + \frac{C_2}{2N_c} \frac{1}{T} \quad (29)$$

for the singlet free energy at infinite separation, and

$$F_1(r, 0) = -\frac{(N_c^2 - 1)}{2N_c} \frac{g^2}{4\pi} \frac{1}{r} + \frac{C_2}{2N_c} \hat{m}_D r \quad (30)$$

for the free energy at zero temperature (i.e., $V_{q\bar{q}}(r)$)⁵. The first thing to observe is that the latter takes the standard zero temperature form of a perturbative Coulombian term plus a linearly raising confining term σr , with

string tension⁶

$$\sigma = \frac{1}{2N_c} C_2 \hat{m}_D. \quad (31)$$

On the other hand, another obvious property in (29) and (30) is the formal resemblance between the two asymptotic free energies with the identification $r \sim 1/T$. More precisely

$$F_1(\infty, T) = F_1(r = 1/m_D, 0). \quad (32)$$

This relation indicates a remarkable and striking duality, within our model, between the quark selfenergy and the zero temperature $q\bar{q}$ potential. Such symmetry can be traced to the form of the propagator $D_{00}(\mathbf{k}; T)$, namely, at least at LO its dependence on momentum and temperature comes in the combination $\mathbf{k}^2 + m_D^2$, regardless of the value of the coupling constant and the non perturbative condensate. Unfortunately the data are not available in momentum space. This makes difficult to check the scaling of the two point function directly. However, when the momentum $\mathbf{k}$ is traded by the separation r , this symmetry can still be seen as a duality between the two asymptotic quantities $F_1(\infty, T)$ and $F_1(r, 0)$.

Of course the abovementioned duality between the roles played by $\mathbf{k}$ and m_D will be disturbed by any scale dependence in the coupling constant and in the condensate corresponding to scaling violations. Such scale dependence implies a possible r and T dependence in those quantities and will be studied further below. Here we take g and C_2 as constant (scale independent) but admit for them the possibility of different values in the two asymptotic regimes ($r, T = 0$) and ($r = \infty, T$). For the coupling constant this assumption is sustained by the data: the bosonic string model fits extremely well the zero temperature $q\bar{q}$ potential, with an error at the level of the 1% in the interval $0.15 \text{ fm} \leq r \leq 0.8 \text{ fm}$ [11]. In this model the coupling constant takes a fixed value

$$g(r, 0) = g_r := \frac{\pi}{2}. \quad (33)$$

Of course asymptotic freedom requires $g(r, 0)$ to go to zero as r decreases, however this is only visible for separations below 0.1 fm [42].⁷ On the other hand, in the large separation regime ($r = \infty, T$), the Polyakov loop data favor a constant g (i.e. a constant a in (11)) rather than the perturbative running one. The constancy of a is unveiled only after the power correction has been subtracted off. Once again, asymptotic freedom requires

⁵ Although the model is devised for the deconfined phase, nothing prevents us from taking the formal zero temperature limit in our expression for the free energy.

⁶ At finite temperature, the $\mathcal{O}(r)$ piece of the free energy is proportional to $m_G^2 - m_D^2$ indicating a weakening of the string tension. Also it confirms that the size of the contact between the zero and finite temperature curves is of the order of $1/m_D$.

⁷ Actually, the coupling $g(\mu) = \pi/2$ matching the string bosonic model corresponds to a NLO evolution scale $1/\mu \sim 0.1 \text{ fm}$ for $\Lambda_{\text{QCD}} \sim 240 \text{ MeV}$.

$g(\infty, T)$ to decrease as T increases but this perturbative regime occurs at very high temperatures only. This “freezing” of $g(\infty, T)$ to a constant value at intermediate temperatures is more thoroughly analyzed from the $q\bar{q}$ potential lattice data in the next subsection. So we take

$$g(\infty, T) = g_T \quad (34)$$

with g_T obtained from the relation

$$a = -\frac{N_c^2 - 1}{2N_c} \frac{g^2}{4\pi} \hat{m}_D. \quad (35)$$

For $N_c = 3$, $N_f = 0$ and $a = -0.19$ this gives

$$g_T = 1.21(11). \quad (36)$$

The fact that both g_r and g_T become frozen at intermediate scales in the two regimes is in support of the duality discussed above.

Let us now turn to the condensate, to see whether the relation (31) between the string tension and the condensate is verified. From the gluodynamics value of b in (12) one obtains a condensate

$$\mathcal{C}_2 = (0.84(4) \text{ GeV})^2. \quad (37)$$

When this is combined in (31) with $g = g_r$ (needed in $\hat{m}_D$), it yields

$$\sigma = (0.43(2) \text{ GeV})^2. \quad (38)$$

Remarkably, this is consistent with the accepted value of the string tension $\sigma \approx (0.42 \text{ GeV})^2$ extracted, e.g., from the slope of the Regge trajectories [43]. In view of this, we will assume that the non perturbative condensate takes a common value in the two asymptotic regimes ($r, T = 0$) and ($r = \infty, T$).

Taking into account that g , but not the condensate, takes two different values in the two regimes, the duality relation (32) gets modified since one has to use g equal to g_T in (29) and equal to g_r in (30).⁸ This yields the refined duality relation

$$F_1(\infty, T) = \gamma^{-3} F_1(r = \gamma/m_D, 0) \quad (39)$$

where

$$\gamma = \left(\frac{g_r}{g_T} \right)^{1/2}, \quad m_D = g_T (N_c/3 + N_f/6)^{1/2} T. \quad (40)$$

To see how this relation works, we plot in Fig. 2 lattice data points available for the two quantities: on the one hand twice the quark selfenergy, $F_1(\infty, T)$ from [10], and on the other the zero temperature $q\bar{q}$ potential $F_1(r, 0)$ from [11]. To compare both quantities we use the relation $r \leftrightarrow \gamma/m_D$ and the rescaling factor γ^3 indicated




FIG. 2: Verification of scaling relations in (39). Black and white squares: $\gamma^3 F_1(\infty, T)$ from lattice data in [10]. Circles: lattice data for the zero temperature $q\bar{q}$ potential from [11]. Both quantities are plotted as functions of the separation r using the relation $r \leftrightarrow \gamma/m_D$.

in (39). γ is computed using the numerical values of g_r and g_T quoted in (33) and (36), which correspond to a continuum extrapolation of those parameters.⁹ The remarkable agreement (modulo simple rescaling) between the two seemingly unrelated quantities suggests that this duality might actually be a feature of QCD, uncovered by our model. (Of course, any duality should break down at temperatures below the transition, since nothing is expected to happen to the zero temperature $q\bar{q}$ potential for separations above $1/T_c$.) The same data are shown in Fig. 3, using the dimensionless quantities $F_1(\infty, T)/T$ and (scaled) $rF_1(r, 0)$, as functions of $(T_c/T)^2$. The straight line patterns (29) and (30) are clearly exhibited, indicating the dominant role played by a non perturbative dimension two operator in the data.

B. Singlet free energy at finite r and T

In order to further check our model against non perturbative information we will use the high quality lattice data of [42], where the singlet $q\bar{q}$ free energy is computed for quenched QCD ($N_c = 3$) at various temperatures in the confining phase for temporal sizes $N_\tau = 4, 8$.

In order to make a comparison we use the following parameterization of the potential

$$V_1(r, T) = - \left(\frac{g^2}{3\pi} \frac{1}{r} + \frac{\mathcal{C}_2}{2N_c T} \right) e^{-\hat{m}_D T r} \quad (41)$$

⁸ Including the g in $\hat{m}_D$, but of course not that in $\mathcal{C}_2$.

⁹ Note that the lattice data for $F_1(r, 0)$, unlike those of $F_1(\infty, T)$, have been extrapolated to the continuum.




FIG. 3: Same lattice data as in Fig. 2 for the dimensionless quantities $F_1(\infty, T)/T$ (squares) and $BrV_{q\bar{q}}(r)$ (circles), as functions of $(T_c/T)^2$. ($B = \hat{m}_D \gamma^{-4}$.) Both the duality and the straight line pattern are exhibited by the data.




FIG. 4: Points: lattice data, for $N_f = 8$, for the singlet free energy as a function of rT , at $T/T_c = 1.05, 1.5, 3, 6.06, 9$, and 12 (from top to bottom, respectively) from Ref. [42]. Curves: $F_1(r, T) = V_1(r, T) + F_1(\infty, T)$ of our model. $g(r)$ (to be used in $V_1(r, T)$) is extracted from the zero temperature potential [11]. In addition $C_2 = (0.9 \text{ GeV})^2$, and $g_T = 1.26$.

where $\hat{m}_D$ is related to g as in (4) with $N_c = 3$, $N_f = 0$. Furthermore, we take an r -dependent g consistent with the zero temperature $q\bar{q}$ -potential in [11], namely, for each r , $g(r)$ is taken so that

$$V_{q\bar{q}}(r) = -\frac{g^2(r)}{3\pi r} + \sigma r. \quad (42)$$

Following [11], for r below 0.1 fm the perturbative value of $V_{q\bar{q}}(r)$ to three-loops is used, and above 0.1 fm , $g(r) = \pi/2$ from the bosonic string model. As noted before this hybrid model reproduces remarkably well the lat-




FIG. 5: Points: values of g_T extracted from a fit to the Polyakov loop data [10, 44] (circles) and from $F_1(r, T)$ lattice data [42] (squares). LO formulas have been used and $C_2 = (0.90(5) \text{ GeV})^2$. The curve displays the perturbative running coupling constant at three-loops.

tice data. The results are presented in Fig. 4 where the model is compared with $N_f = 8$ lattice data (the results for $N_f = 4$ data are qualitatively similar). For clarity we represent F_1 instead of V_1 , and use rT as the independent variable. For the temperature dependent shift $F_1(\infty, T)$ the expression (29) is used with $g_T = 1.26$, which corresponds to a value $a = -0.21$; this is the LO perturbative value around $T = 6 T_c$. For the condensate we take $C_2 = (0.9 \text{ GeV})^2$ which gives a fair account of $F_1(\infty, T)$ at lower temperatures. (Note that the points in the figure are the actual lattice data, without any continuum extrapolation.) For consistency the same value of C_2 is used in $V_1(r, T)$ but the precise value is not very important in the potential (also it is not very important for $F_1(\infty, T)$ for high temperatures). As we can see from Fig. 4, the overall pattern of the lattice data is reasonably well reproduced by our model. Given the small error bars in the data, a more detailed agreement (small χ^2/pdof) was not to be expected without further sophistication of the model. As was noted in Section II, $F_1(\infty, T)$ (or equivalently, the Polyakov loop) cannot be described at all without introducing a dimension two condensate. The same holds here for the potential: the overall shape of lattice data for $V_1(r, T)$ cannot be reproduced even approximately if C_2 is set to zero, moreover, the same value of the condensate extracted from $F_1(\infty, T)$ works for the potential too. This feature is expected to survive in more elaborated models.

We have considered also a fit to the data allowing a T dependent g_T (but keeping C_2 constant). This is depicted in Fig. 5. The fit using all data for the free energy is of course more stable than that using only the Polyakov loop, although both sets yield consistent results. The error bars are very small at high temperatures and come

from the lattice data and from allowing a variation range $C_2 = (0.90 \pm 0.05 \text{ GeV})^2$ in the condensate. We find an almost flat dependence of g_T with the temperature, much flatter than expected from a perturbative running. As noted, a nearly constant g_T is in favor of the duality relation displayed in Fig. 2 and 3, since g_r is also frozen in the bosonic string model. g_T is somewhat larger than the perturbative value. It is noteworthy that a larger renormalization of the coupling constant, of about 50%, has also been observed in [42, 45] if the observed Debye mass, extracted from the exponential falloff of the potential, is compared to its perturbative value. In our model the exponential fall at large separations is controlled by g_r rather than g_T . Since $g_r = \pi/2$ is rather large, as compared to the perturbative value, this is qualitative agreement with the findings in [42, 45] (see also [46]), however, g_r is somewhat smaller and does not run.

The present analysis of the singlet $q\bar{q}$ free energy is carried out at LO, and further studies will have to take into account higher orders in the loop expansion. We certainly do not expect these higher order corrections to reproduce the effect of condensates. In addition, corrections coming from the condensate anomalous dimension are also expected to be relevant in more accurate descriptions; at the present stage this effect on the non-perturbative side should play a comparable role to loops corrections. Finally, higher dimensional condensates could also be present and affect the free energy. Unfortunately, as our previous analysis reveals [22], the extraction of a dimension four condensate from the lattice data of the Polyakov loop is not reliable at present. It would contribute to (11) with a term $(T_c/T)^4$.

V. CONCLUSIONS

In the present paper we have investigated the role of dimension two condensates in the interrelation between the Polyakov loop at finite temperature above the phase transition and the static quark-antiquark potential at zero temperature. This extends our previous model for the renormalized Polyakov loop in the deconfined phase which proved very successful and enabled a quantitative determination of the BRST invariant dimension two

gluon condensate [22]. Here we have studied the consequences of the model when applied to the heavy quark-antiquark free energy in the unconfined phase of QCD.

In our approach the perturbative and non-perturbative contributions are explicitly separated and an analytic and simple expression for the free energy which interpolates between the short and long distances regime is derived. The separation turns out to accommodate quite naturally a smooth running coupling when the model is confronted to lattice data, which speaks in favor of the consistency of the approach.

We observe a natural consequence of the presence of the dimension two condensate if one restricts to LO in the perturbative contributions and effects associated to anomalous dimensions are disregarded. Although in general the free energy of a quark-antiquark pair depends both on the distance r and the temperature T , there are some regions where this double dependence presents strong correlations. Amazingly, the regimes of finite temperature and long distances (from which the Polyakov loop is defined) and the zero temperature and finite distances (from which the zero temperature $q\bar{q}$ potential is obtained) have a striking dual functional dependence which is verified on a quantitative level on the lattice if the simple change of variable $r \leftrightarrow 1/m_D$ is assumed. The very fact that the non-perturbative contributions are encoded in the dimension two condensate suggests that our result holds up to mild logarithmic radiative corrections and it would be interesting to quantify the observed small discrepancies in a more systematic manner. Despite all these caveats we believe that the quantitative success of the duality relations provides a further evidence on existence and relevance of dimension two gluon condensates.

Acknowledgments

We thank O. Kaczmarek and F. Karsch for kindly providing us with the lattice data of Ref. [42]. This work is supported in part by funds provided by the Spanish DGI and FEDER funds with grant no. FIS2005-00810, Junta de Andalucía grants no. FQM225-05 and FQM481 and EU Integrated Infrastructure Initiative Hadron Physics Project contract no. RII3-CT-2004-506078.

-
- [1] L. D. McLerran, Rev. Mod. Phys. **58**, 1021 (1986).
 - [2] J. Adams et al. (STAR), Nucl. Phys. **A757**, 102 (2005), nucl-ex/0501009.
 - [3] R. M. Weiner, Int. J. Mod. Phys. **E15**, 37 (2006), hep-ph/0507115.
 - [4] A. Mocsy and P. Petreczky, Eur. Phys. J. **C43**, 77 (2005), hep-ph/0411262.
 - [5] W. M. Alberico, A. Beraudo, A. De Pace, and A. Molinari (2006), hep-ph/0612062.
 - [6] T. Matsui and H. Satz, Phys. Lett. **B178**, 416 (1986).
 - [7] C. P. Korthals Altes, Nucl. Phys. **B420**, 637 (1994), hep-th/9310195.
 - [8] R. D. Pisarski (2002), hep-ph/0203271.
 - [9] E. Gava and R. Jengo, Phys. Lett. **B105**, 285 (1981).
 - [10] O. Kaczmarek, F. Karsch, P. Petreczky, and F. Zantow, Phys. Lett. **B543**, 41 (2002), hep-lat/0207002.
 - [11] S. Necco and R. Sommer, Nucl. Phys. **B622**, 328 (2002), hep-lat/0108008.
 - [12] F. A. Chishtie and V. Elias, Phys. Lett. **B521**, 434 (2001), hep-ph/0107052.
 - [13] A. Pineda, JHEP **06**, 022 (2001), hep-ph/0105008.
 - [14] G. Cvetic, J. Phys. **G30**, 863 (2004), hep-ph/0309262.

- [15] A. H. Hoang, M. C. Smith, T. Stelzer, and S. Willenbrock, Phys. Rev. **D59**, 114014 (1999), hep-ph/9804227.
- [16] M. Beneke, Phys. Lett. **B434**, 115 (1998), hep-ph/9804241.
- [17] N. Brambilla, A. Pineda, J. Soto, and A. Vairo, Phys. Rev. **D60**, 091502 (1999), hep-ph/9903355.
- [18] G. S. Bali, Phys. Rept. **343**, 1 (2001), hep-ph/0001312.
- [19] Y. Sumino, Phys. Lett. **B571**, 173 (2003), hep-ph/0303120.
- [20] O. Andreev and V. I. Zakharov (2006), hep-ph/0611304.
- [21] K. G. Wilson, Phys. Rev. **179**, 1499 (1969).
- [22] E. Megías, E. Ruiz Arriola, and L. L. Salcedo, JHEP **01**, 073 (2006), hep-ph/0505215.
- [23] K.-I. Kondo, Phys. Lett. **B514**, 335 (2001), hep-th/0105299.
- [24] E. Megías, E. R. Arriola, and L. L. Salcedo, Rom. Rep. Phys. **58**, 081 (2006), hep-ph/0510114.
- [25] S. Nadkarni, Phys. Rev. **D27**, 917 (1983).
- [26] P. Ginsparg, Nucl. Phys. **B170**, 388 (1980).
- [27] T. Appelquist and R. D. Pisarski, Phys. Rev. **D23**, 2305 (1981).
- [28] E. Braaten and A. Nieto, Phys. Rev. **D53**, 3421 (1996), hep-ph/9510408.
- [29] M. E. Shaposhnikov (1996), effective theories and fundamental interactions, Erice 1996, 360-383, hep-ph/9610247.
- [30] E. Megías, E. R. Arriola, and L. L. Salcedo (2006), hep-ph/0610163.
- [31] A. Dumitru, Y. Hatta, J. Lenaghan, K. Orginos, and R. D. Pisarski, Phys. Rev. **D70**, 034511 (2004), hep-th/0311223.
- [32] O. Kaczmarek and F. Zantow, Phys. Rev. **D71**, 114510 (2005), hep-lat/0503017.
- [33] P. Pascual and R. Tarrach, Lect. Notes Phys. **194**, 1 (1984).
- [34] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. **B147**, 385 (1979).
- [35] S. Narison, World Sci. Lect. Notes Phys. **26**, 1 (1989).
- [36] M. J. Lavelle and M. Schaden, Phys. Lett. **B208**, 297 (1988).
- [37] K. G. Chetyrkin, S. Narison, and V. I. Zakharov, Nucl. Phys. **B550**, 353 (1999), hep-ph/9811275.
- [38] P. Boucaud et al., Phys. Rev. **D63**, 114003 (2001), hep-ph/0101302.
- [39] E. Ruiz Arriola, P. O. Bowman, and W. Broniowski, Phys. Rev. **D70**, 097505 (2004), hep-ph/0408309.
- [40] P. Boucaud et al. (2005), hep-lat/0504017.
- [41] O. Philipsen, Phys. Lett. **B535**, 138 (2002), hep-lat/0203018.
- [42] O. Kaczmarek, F. Karsch, F. Zantow, and P. Petreczky, Phys. Rev. **D70**, 074505 (2004), hep-lat/0406036.
- [43] A. V. Anisovich, V. V. Anisovich, and A. V. Sarantsev, Phys. Rev. **D62**, 051502 (2000), hep-ph/0003113.
- [44] F. Zantow (2003), hep-lat/0301014.
- [45] O. Kaczmarek and F. Zantow (2005), hep-lat/0512031.
- [46] N. O. Agasian and Y. A. Simonov, Phys. Lett. **B639**, 82 (2006), hep-ph/0604004.